



Lancaster Plastic Surgery Associates
1496 Lititz Pike • Lancaster, PA 17601 • (717) 481-3444

Post-Procedure Instructions

Endoscopic Carpal Tunnel Release

You had an endoscopic carpal tunnel release with one or two small (approximately 1 cm) incisions closed with dissolvable buried sutures and skin glue (Dermabond). Please follow these instructions carefully.

Dressing & Wound Care

- Leave the initial dressing in place for 2–3 days.
- After 2–3 days you may remove the outer dressing.
- The skin glue (Dermabond) will protect the incision. It will gradually peel off on its own over 7–14 days—do not pick at it.
- You may gently wash the area with soap and water once the dressing is off. Pat dry—do not rub.
- Avoid soaking the hand (no baths, pools, or hot tubs) for at least 1 week or until the glue has fully come off.
- A light bandage may be used if desired for comfort.

Elevation & Swelling

- Keep your hand elevated above the level of your heart as much as possible for the first 48–72 hours.
- This reduces swelling and discomfort.

Activity & Restrictions

- Move your fingers gently and frequently to help reduce stiffness.
- Avoid heavy lifting, gripping, or repetitive forceful use of the hand for at least 2 weeks (or as directed).
- Most patients can resume light daily activities within a few days.
- Driving is usually safe once you feel comfortable and are no longer taking narcotic pain medication.

Pain Management

- Mild to moderate discomfort is normal for several days.
- Take over-the-counter acetaminophen or ibuprofen as directed unless told otherwise.
- If stronger medication was prescribed, use it only as needed.

Follow-Up Visits

- Two post-operative visits are included in your package.
- Please keep your scheduled appointments so we can check healing progress.

Scar Care

- Beginning at 2 weeks, you may start gentle scar massage.
- Massage the small scars with a moisturizing lotion or silicone gel for a few minutes several times a day.
- Scar management guidance is part of your package.

When to Call the Office

- Increasing redness, warmth, swelling, or drainage from the incision
- Fever over 101°F (38.3°C)
- Severe or worsening pain not controlled by medication
- New or increasing numbness or weakness
- Any other concern that worries you

Office: (717) 481-3444

If you have an urgent concern after hours, call the office number for instructions.