



Immunization Schedule for Midland Pediatric Associates

The following immunization schedule is the routine childhood immunization schedule that we use at Midland Pediatric Associates and fully complies with the immunization schedule as required by the Texas state law and as recommended by the ACIP/AAP/AAFP*.

*ACIP – Advisory Committee on Immunization Practices | *AAP – American Academy of Pediatrics | *AAFP – American Academy of Family Physicians

REQUIRED IMMUNIZATIONS

Age	Immunization	Method
Birth	HBV	1 injection
2 months	(DTaP + IPV + HBV + HIB), PCV, Rotavirus	2 injections, 1 oral
4 months	(DTaP + IPV + HBV + HIB), PCV, Rotavirus	2 injections, 1 oral
6 months	(DTaP + IPV + HBV + HIB), PCV, Rotavirus	2 injections, 1 oral
12 months	HIB, HAV, PCV	3 injections
15 months	MMR, Varicella, DTaP	3 injections
2 years	HAV (may be given at 18 month visit)	1 injection
4 years	MMR, (DTaP + IPV), Varicella	3 injections
11 years	Tdap, MCV4	3 injections
16-18 years	MCV4	1 injection

RECOMMENDED IMMUNIZATIONS

Age	Immunization	Dose #	Interval
6 mos-8 years	Influenza (IIV4-Injection)	2 doses	≥ 4 weeks apart (1 st year)
≥ 2 years	Influenza (LAIV4 - Nasal)	1 dose	Annually
≥ 9 years	Influenza (IIV4 -Injection)	1 dose	Annually
≥ 5 years	COVID-19	2 doses	3 weeks apart
≥ 9 years	HPV	2 doses	6 months apart
≥ 16 years	MenB	2 doses	1 month apart

HAV Hepatitis A
 HBV Hepatitis B
 HIB Haemophilus Influenza Type B
 DTaP Diphtheria, Tetanus, Acellular Pertussis
 IPV (Injectable) Polio
 Varicella Chicken Pox
 MMR Measles, Mumps, Rubella
 PCV Pneumococcal Conjugate
 HAV Hepatitis A
 Rotavirus Rotavirus
 Tdap Tetanus, Diphtheria, Acellular Pertussis
 MCV4 Meningococcal Conjugate (serogroups ACWY)

Influenza IIV4 - Inactivated Quadrivalent Injection
 6 mos-8 yrs - (2 doses if have NOT previously received ≥ 2 doses of trivalent or quadrivalent influenza vaccine in the same influenza season – only 1 dose if have been boosted in previous influenza season)
 Influenza LAIV4 - Live Attenuated Nasal Inhalation
 COVID-19 SARS-CoV-2 (Coronavirus/COVID-19) (2 dose primary series approved for ages ≥ 5 – booster optional for ≥ age 12 at ≥ 5 months after 2 dose in primary series)
 HPV Human Papilloma Virus (2 dose series < age 15 [3 dose series > age 15, 1st dose at 0, 2nd dose at 1 month, 3rd at 6 months] approved for ages 9-26)
 MenB Meningococcal B (2 dose series – 1 month apart)