



CornSpike

Premium Liquid Fertilizer Solution

GUARANTEED ANALYSIS

Boron (B)	0.1%
Copper (Cu)	0.5%
0.5% Chelated Copper (Cu)	
Iron (Fe)	0.25%
0.25% Chelated Iron (Fe)	
Manganese (Mn)	1.0%
Zinc (Zn)	5.0%
5.0% Chelated Zinc (Zn)	

Derived From Borid Acid, Copper PASP, Iron EDTA
Iron PASP, Manganese EDTA, Manganese PASP,
Zinc EDTA and Zinc PASP

WARNING: The use of Boron on any crops other than those recommended may result in injury to the crop. Contact your county agent or field representative before making application.

CAUTION: KEEP OUT OF REACH OF CHILDREN



WARNING

Harmful if swallowed. Harmful in contact with skin. Causes skin irritation. Causes eye irritation. Harmful if inhaled. May cause respiratory irritation. May cause drowsiness or dizziness. Suspected of damaging fertility or the unborn child.

PRECAUTIONARY STATEMENTS:

Do not handle until all safety precautions have been read and understood. Avoid breathing fume/mist/vapours/spray. Wash skin thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear protective gloves/protective clothing/eye and face protection.

IF SWALLOWED: Call a poison control center or doctor/physician if you feel unwell. Rinse mouth.

IF ON SKIN: Wash with plenty of soap and water. Call a poison center or doctor/physician if you feel unwell. If skin irritation persists: Get medical advice/attention.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Call a poison control center or doctor/physician if you feel unwell.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do - continue rinsing. If eye irritation persists: Get medical advice/attention.

Take off contaminated clothing and wash it before reuse. Store in a well-ventilated place. Keep container tightly closed. Store locked up. Dispose of contents/container to an approved waste disposal plant in accordance with local/regional/national regulations.

Application Instructions

QLF CornSpike can be applied to all types of crops.

Maintenance Rate: Apply 1 to 2 quarts per acre.

Medium to Severe Deficiency Rate: Apply 3 to 4 quarts per acre.

Applications should be made at or prior to planting or at the start of the growing season. Applications should be made every 7 - 28 days as needed throughout the season.

Conditions of Sale

Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions and use. No one other than an officer of the seller is authorized to make any warranty, guarantee, or direction concerning this product. Because time, place, rate of application and other conditions of use are beyond the seller's control, the seller's liability from handling, storage, and use of this product is limited to replacement of the product or a refund of the purchase price.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm> or by writing to QLF at the address on this label and referring to the lot/batch # on this container.

Weight per Gallon: 10.7 lbs (4.8 kg) @ 68° F