

H-85™ Flowable

INCREASES MICROBIAL DIVERSITY



Guaranteed Analysis (0-0-5)

Soluble Potash (K₂O)5%

Derived from potassium hydroxide.

ALSO CONTAINS NON-PLANT FOOD INGREDIENT(S)

GUARANTEED ANALYSIS - SOIL AMENDING INGREDIENT(S):

12%..... Humic Acid from Leonardite

88%.....Inert Ingredients (as a non-plant food ingredient)

General Information

H-85™ Flowable is a multi-extraction humic and fulvic acid that contains short-, medium-, and long-chain soluble carbon molecules. **H-85™ Flowable** is soluble across a broad pH range, improving soluble carbon availability in all soil types. The availability of soluble carbon across a broad pH range increases soil microbial diversity and soil health. When combined with synthetic nitrogen, **H-85™ Flowable** will increase nitrogen efficiency by at least 25%.

Product Usage

Water Penetration	Apply 0.125-0.5 gals./acre as a soil application every 1 to 4 weeks.
Soil Health	Apply 0.125-0.5 gals./acre as a soil application every 1 to 4 weeks.
Root Growth	Apply 0.125-0.5 gals./acre as a soil application every 1 to 4 weeks.
Nitrogen Efficiency	Apply 0.125-0.5 gals./acre as a soil application every 1 to 4 weeks.



H-85™ Flowable

INCREASES MICROBIAL DIVERSITY

H-85™ Flowable Handling Guidelines

PREMIXING

- Premixing is considered a best practice when sprayer agitation is not optimum. Proper hydration is essential for all applications. Recirculate or agitate while adding material.

COMPATIBILITY

- Always jar test first.
- Redox products are compatible with other Redox products when following product handling guidelines.
- Use caution with reactive materials, such as phosphorus and calcium.
- Avoid extreme shifts in tank pH. When utilizing Redox materials that acidify, check tank pH prior to adding buffers.

TANK MIXING

- Use of an anti-foaming agent is recommended. Fill the tank 50% full with water and initiate tank agitation prior to adding materials.
- Don't add material too quickly—this allows for more thorough hydration.
- The use of inductor assemblies is encouraged.
- Recirculate or agitate while adding material.
- Total foliar spray content of dry soluble Redox product should not exceed 2% solution (by weight).
- If material is not applied immediately, tank recirculation is required prior to application to ensure uniform product distribution.

REFER TO PRODUCT HANDLING GUIDELINES FOR ADDITIONAL MIXING INSTRUCTIONS.



Scan QR code for
product information