


Mainstay™

Calcium 2.0

IMPROVES CALCIUM NUTRITION &
BUILDS SOIL STRUCTURE



Guaranteed Analysis

Calcium (Ca)..... 20%
Derived from calcium carbonate

Organic version available: 

General Information

Mainstay™ Calcium 2.0 is specifically formulated to provide concentrated plant-available calcium for plant nutrition and soil structure.

CALCIUM NUTRITION

Redox microencapsulation allows for improved calcium uptake and movement within the plant.

SOIL STRUCTURE

Redox microencapsulation prevents calcium tie-up in the soil, allowing for improved soil structure.

Product Usage

Fruit Firmness	Apply 0.5-5 gals./acre as a soil application or 0.25-1 gal./acre as a foliar application during early season cell division/development. Repeat every 1 to 2 weeks.
Cell Wall Strength	Apply 0.5-5 gals./acre as a soil application or 0.25-1 gal./acre as a foliar application during early season cell division/development. Repeat every 1 to 2 weeks.
Plant Nutrition	Apply 0.5-5 gals./acre as a soil application or 0.25-1 gal./acre as a foliar application during early season cell division/development. Repeat every 1 to 2 weeks.
Soil Structure	Apply 0.5-5 gals./acre as a soil application as needed. Repeat every 2 to 4 weeks.
Root Growth	Apply 0.5-5 gals./acre as a soil application as needed. Repeat every 2 to 4 weeks.



Mainstay™

Calcium 2.0

IMPROVES CALCIUM NUTRITION &
BUILDS SOIL STRUCTURE

Mainstay™ Calcium 2.0 Handling Guidelines

PREMIXING

- Premixing is considered a best practice when sprayer agitation is not optimum. Proper hydration is essential for all applications. Recirculate or agitate while adding material.

COMPATIBILITY

- Always jar test first.
- Redox products are compatible with other Redox products when following product handling guidelines.
- Use caution with reactive materials, such as phosphorus and calcium.
- Avoid extreme shifts in tank pH. When utilizing Redox materials that acidify, check tank pH prior to adding buffers.

TANK MIXING

- Use of an anti-foaming agent is recommended. Fill the tank 50% full with water and initiate tank agitation prior to adding materials.
- Don't add material too quickly—this allows for more thorough hydration.
- The use of inductor assemblies is encouraged.
- Recirculate or agitate while adding material.
- Total foliar spray content of dry soluble Redox product should not exceed 2% solution (by weight).
- If material is not applied immediately, tank recirculation is required prior to application to ensure uniform product distribution.

REFER TO PRODUCT HANDLING GUIDELINES FOR ADDITIONAL MIXING INSTRUCTIONS.



Scan QR code for
product information