

Rx Platinum™

IMPROVES ROOT GROWTH &
MICROBIAL DIVERSITY



Guaranteed Analysis (1-7-12)

Total Nitrogen (N)	1%
0.25%.....	Ammoniacal Nitrogen
0.25%.....	Nitrate Nitrogen
0.50%.....	Urea Nitrogen
Available Phosphate (P ₂ O ₅)	7%
Soluble Potash (K ₂ O)	12%

Derived from ammonia nitrate, urea, phosphoric acid and potassium hydroxide.

ALSO CONTAINS NON-PLANT FOOD INGREDIENT(S)

GUARANTEED ANALYSIS - SOIL AMENDING INGREDIENT(S):

2%.....	Humic Acid from Leonardite
98%.....	Inert Ingredients (as a non-plant food ingredient)

General Information

Rx Platinum™ is a phosphorus and potassium fertilizer, containing proprietary soluble carbon compounds. Rx Platinum™ reduces abiotic stress, improves root growth, and increases soil microbial diversity due to humic, fulvic, and long-chain soluble carbon content. Rx Platinum™ utilizes technology from its proprietary formulations of diKaP™, H-85™, and Rootex™.

Product Usage

Abiotic Stress Defense	Apply 0.5 -3 gals./acre as a foliar/soil application every 1 - 6 weeks. Water in.
Heat Stress	Apply 0.5 - 3 gals./acre as a foliar/soil application every 1 - 6 weeks. Water in.
Root Growth	Apply 0.5 - 3 gals./acre as a soil application every 1 - 6 weeks. Water in.
Soil Health	Apply 1 - 3 gals./acre as a foliar/soil application every 1 - 6 weeks. Water in.



Rx Platinum™

IMPROVES ROOT GROWTH &
MICROBIAL DIVERSITY

Rx Platinum™ Handling Guidelines

PREMIXING

- Premixing is considered a best practice when sprayer agitation is not optimum. Proper hydration is essential for all applications. Recirculate or agitate while adding material.

COMPATIBILITY

- Always jar test first.
- Redox products are compatible with other Redox products when following product handling guidelines.
- Use caution with reactive materials, such as phosphorus and calcium.
- Avoid extreme shifts in tank pH. When utilizing Redox materials that acidify, check tank pH prior to adding buffers.

TANK MIXING

- Use of an anti-foaming agent is recommended. Fill the tank 50% full with water and initiate tank agitation prior to adding materials.
- Don't add material too quickly—this allows for more thorough hydration.
- The use of inductor assemblies is encouraged.
- Recirculate or agitate while adding material.
- Total foliar spray content of dry soluble Redox product should not exceed 2% solution (by weight).
- If material is not applied immediately, tank recirculation is required prior to application to ensure uniform product distribution.

REFER TO PRODUCT HANDLING GUIDELINES FOR ADDITIONAL MIXING INSTRUCTIONS.



Scan QR code for
product information