

TriPlex™

# MICRO Flowable

IMPROVES MICRONUTRIENT NUTRITION & NITROGEN METABOLISM



## Guaranteed Analysis

Boron (B).....	0.66%
Copper (Cu).....	0.66%
Iron (Fe).....	2.1%
Manganese (Mn).....	2.1%
2.1%.....	Water Soluble Manganese (Mn)
Zinc (Zn).....	2.1%

Derived from boric acid, copper sulfate, ferrous sulfate, manganese sulfate, and zinc sulfate.  
Warning Boron: The use of the fertilizing material containing boron on any crops other than those recommended may result in serious injury to the crop(s).

## General Information

TriPlex™ Micro Flowable is an L-amino acid micro-nutrient fertilizer. Redox chelation and complexing technology increases micro-nutrient availability. TriPlex™ Micro Flowable is designed with specific micronutrient ratios that improve nitrogen metabolism.

## Product Usage

Plant Nutrition	Apply 0.125-1 gal./acre as a soil application every 1 to 4 weeks, or 0.125-1 gal./acre as a foliar application every, 1 to 4 weeks. Water in.
Rapid Growth & Cell Expansion	Apply 0.06-1.5 gal./acre as a foliar/soil application 1 to 4 weeks.
Root Growth	Apply 0.06-0.5 gal./acre as a foliar/soil application 1 to 4 weeks.
Crop Set & Sizing	Apply 0.06-0.5 gal./acre as a foliar/soil application 1 to 4 weeks.



TriPlex™

# MICRO Flowable

IMPROVES MICRONUTRIENT NUTRITION &  
NITROGEN METABOLISM

---

## TriPlex™ Micro Flowable Handling Guidelines

### PREMIXING

- Premixing is considered a best practice when sprayer agitation is not optimum. Proper hydration is essential for all applications. Recirculate or agitate while adding material.

### COMPATIBILITY

- Always jar test first.
- Redox products are compatible with other Redox products when following product handling guidelines.
- Use caution with reactive materials, such as phosphorus and calcium.
- Avoid extreme shifts in tank pH. When utilizing Redox materials that acidify, check tank pH prior to adding buffers.

### TANK MIXING

- Use of an anti-foaming agent is recommended. Fill the tank 50% full with water and initiate tank agitation prior to adding materials.
- Don't add material too quickly—this allows for more thorough hydration.
- The use of inductor assemblies is encouraged.
- Recirculate or agitate while adding material.
- Total foliar spray content of dry soluble Redox product should not exceed 2% solution (by weight).
- If material is not applied immediately, tank recirculation is required prior to application to ensure uniform product distribution.

*REFER TO PRODUCT HANDLING GUIDELINES FOR ADDITIONAL MIXING INSTRUCTIONS.*

---



Scan QR code for  
product information