

NUTRIENTS SUPPLIED (pounds per gallon):

0.204
0.204
0.010
0.015
0.031
0.153
0.00005
0.409

Derived From: Ammonium Hydroxide, Potassium Hydroxide, Boric Acid, Copper EDTA, Iron EDTA, Manganese EDTA, Sodium Molybdate, Zinc EDTA

PRODUCT PROPERTIES:

Analysis:	2-0-21B15Cu3Fe-
	1.5Mn0005Mo-4Zn
Weight:	10.22 lbs. per gallon
Specific gravity:	1.225 kg/L
pH:	6.8-7.5
Appearance:	Clear, brown liquid
Odor:	Slight amine

GENERAL PRODUCT INFORMATION:

NACHURS CropMax liquid fertilizer is manufactured with 100% fully EDTA chelated copper, manganese, iron, and zinc. Unlike other micronutrient sources such as complexes, partial chelates, and natural organic complexes, NACHURS EDTA chelated micronutrients are 100% available to the crop. Other micro sources contain too little complexing agent and undergo major chemical changes, delivering significantly less micronutrient in a form available for plant uptake. While these sources of micros may offer cost savings at first, they can actually create deficiencies for lack of availability.

FIRST AID: Please see SDS sheet for more information, call (800) 622-4877 or visit us online at www.nachurs.com.

KEEP OUT OF REACH OF CHILDREN.

*These are general product recommendations. Please consult with your authorized NACHURS distributor or agronomist for specific fertility recommendations. These recommendations are believed to be reliable and should be followed carefully. Failure to follow label directions or improper application practices, all of which are out of control of the manufacturer or seller, can result in plant or leaf damage. Crop injury may result from unusual weather conditions, failure to follow label directions or improper application practices all of which are out of control of NACHURS.

SELLER WARRANTS THAT THE ABOVE PRODUCT CONFORMS TO ITS CHEMICAL DESCRIPTION AND IS REASONABLY FIT FOR THE PURPOSE ON THE LABEL WHEN USED IN ACCORDANCE WITH DIRECTIONS UNDER NORMAL CONDITIONS OF USE (INCLUDING NORMAL WEATHER CONDITIONS). NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXPRESS OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT WHEN USED CONTRARY TO THE LABEL INSTRUCTIONS OR UNDER ABNORMAL CONDITIONS (INCLUDING ABNORMAL WEATHER CONDITIONS), AND THE BUYER ASSUMES THE RISK OF ANY SUCH USE. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS

© 2022 Nachurs Alpine Solutions. All rights reserved.

NACHURS CropMax®

Premium Liquid Micronutrients



APPLICATION RATES:

• In-furrow - 1-2 pints per acre.

NACHURS CropMax is formulated to provide necessary micronutrients for optimum growth with the advantage of excellent compatibility with NACHURS in-furrow

- Foliar apply 1–2 pints per acre with sufficient spray volume to ensure good coverage.
- Fertigation apply 1–2 pints per acre. Extremely hard water and/or high total dissolved solids may effect performance.

Compatibility: ${\bf NACHURS\ CropMax\ }$ may be applied in combination with other liquid fertilizers, fertilizer suspensions, and nitrogen solutions. Always jar test combinations before field mixing.

These are general product recommendations, Please consult with your NACHURS Sales Manager or agronomist for specific fertility recommendations.

GENERAL MIXING INSTRUCTIONS

- 1. Put 1/3 of fertilizer in tank
- 2. Add other chemicals, if any
- 3. Fill tank with balance of fertilizer
- 4. Add correct amount of chelated miccronutrient
- 5. Agitate adequately to mix

CAUTION: Check compatibility with standard jar test.

WARNING: BORON IS TO BE USED WHERE SOIL TEST AND/OR TISSUE ANALYSIS INDICATE A DEFICIENCY, AND SHOULD NOT BE USED AT RATES IN EXCESS OF THE RATE RECOMMENDED BY A QUALIFIED INDIVIDUAL/ENTITY SUCH AS A CERTIFIED CROP ADVISOR, AGRONOMIST OR UNIVERSITY. EXCESSIVE APPLICATION OF BORON MAY CAUSE CROP DAMAGE. THIS FERTILIZER IS TO BE USED ONLY ON SOIL WHICH RESPONDS TO MOLYBDENUM. CROPS HIGH IN MOLYBDENUM ARE TOXIC TO GRAZING ANIMALS (RUMINANTS).

THE ROLE OF MICRONUTRIENTS:

Boron is vital to the growth and development of the plant. Without adequate boron, new growth ceases. It is necessary in the pollination and seed production stages. Boron is essential for maintaining a balance between sugars and starches. A small amount of boron is beneficial to plants but too much can be toxic to

Copper (Cu)

Copper is important as a co-enzyme. It is needed to activate several plant enzymes, including building and converting amino acids to proteins. Since copper is an immobile nutrient, deficiency symptoms usually occur on new growth. Copper deficient plants will become chlorotic and take on a bleached appearance. New growth may die.

Iron (Fe)

Iron is required for the formation of chlorophyll in plant cells. It serves as an activator for biochemical processes such as respiration, photo-synthesis and symbiotic nitrogen fixation.

Manganese (Mn)

Manganese is essential to plants but too much is toxic. Manganese functions in chlorophyll development and serves as a catalyst in several enzyme systems in the oxidation-reduction process. Manganese deficiencies are very similar to iron deficiencies and appears in the younger leaves of the plant first. Color may be pale between the veins of broadleaf plants.

Molybdenum (Mo)

Molybdenum helps to transform basic nitrogen into amino acids, which are building blocks for proteins. It also helps legumes to symbiotically fix atmospheric nitrogen. Molybdenum is a catalyst in many oxidationreduction process within plants.

Zinc (Zn)

Zinc is necessary for starch formation and proper root development. It is also essential for seed formation and maturity. The most common symptom of zinc deficiency is interveinal chlorosis on older leaves with shortening of the internodal area. This shortening often results in a short compressed plant with a rosetted appearance.



















