

Rooster Booster™

**GUARANTEED
ANALYSIS
0.3-0.2-0.4**

Total Nitrogen (N) 0.3%
 0.2% Ammoniacal Nitrogen
 0.0% Other Water-Soluble Nitrogen
 0.1% Water Insoluble Nitrogen

Available Phosphate (P₂O₅) 0.2%

Total Potassium (K₂O) 0.4%
 0.4% Potassium

Derived from: Chicken Manure

CAUTION:
KEEP OUT OF REACH OF CHILDREN
 Wash hands and face before eating or drinking. Wash soiled clothing before wearing again if contact occurs.
EYES: Contact can cause mild irritation of eyes. Flush with running water. Seek medical attention if irritation persists.
SKIN: Wash thoroughly with soap & water. Seek medical attention if rash develops.
IF SWALLOWED: Drink one or two glasses of water and induce vomiting. Seek immediate attention. Never induce vomiting or give anything by mouth to an unconscious person.
INHALATION: Move to fresh air. If irritation persists, get medical attention.



NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used according to directions under normal conditions on most agricultural crops. Because of widely varying use conditions, it is impossible to eliminate all risks even when the label directions are followed. Therefore, the manufacturer makes no other warranties, expressed or implied. The buyer assumes all risks associated with the use of this product.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Crop Type:	Rate/Ac	Application Type:	Timing:
Grass, Hay, Pasture	3-5 gallons	Foliar	Contact your representative
Corn	1-5 gallons	2x2, In-Furrow, Y-Drop, Foliar	Planting, Sidedress, V3-V6, VT
Legumes	1-5 gallons	In-Furrow, Foliar	Planting, V3-V5, R1-R4
All other row crops	1-5 gallons	In-Furrow, Foliar	Contact your representative
Commercial Produce	3-10 gallons	Fertigation, Foliar	Vegetative growth, fruit/pod set

Rooster Booster can be applied with liquid nitrogen and other foliar applied nutrients. Rooster Booster can be applied with humic and fulvic acid products, as well as bio-stimulants including sugar based products.

Always perform a jar test before mixing large batches.

Distributed by:



Weight per Gallon: 8.5 lbs per gallon at 68° Fahrenheit
 □ 275 Gal (1.0 Kl) Net wt 2,337.5 lbs