



## Fever/Pain Relief Medicines for Infants/Children

\* Check the concentration (MG/ML) and type of medicine on the bottle you are using **BEFORE** dosing your child \*

### Tylenol® (Acetaminophen) dosing chart for home use

· Tylenol® (Acetaminophen) can be given every 4 hours

· For a child with vomiting/seizures who can't take oral Acetaminophen, rectal suppositories are available in 120 mg, 300 mg, and 650 mg  
(Use dosing chart below to find the dose in mg needed per your child's current weight)

Weight in Pounds (lb)	Weight in Kilograms (kg)	Dosage in Mg	Infant Drops 160 mg/5 ml	Children's Liquid 160 mg/5 ml 32 mg/ml	Chewable Tablets 80 mg tabs	Junior Strength Tablets 160 mg tabs	Adult Strength Tablets 325 mg tabs
6 - 11	3 - 5	40 mg	1.25 ml	1/4 teaspoon 1.25 ml			
12 - 17	6 - 7	80 mg	2.5 ml	1/2 teaspoon 2.5 ml			
18 - 23	8 - 10	120 mg	3.75 ml	3/4 teaspoon 3.75 ml			
24 - 35	11 - 15	160 mg	5 ml	1 teaspoon 5 ml	2 tablets	1 tablet	
36 - 47	16 - 21	240 mg		1 1/2 teaspoons 7.5 ml	3 tablets	1 1/2 tablets	
48 - 59	22 - 26	320 mg		2 teaspoons 10 ml	4 tablets	2 tablets	1 tablet
60 - 71	27 - 32	400 mg		2 1/2 teaspoons 12.5 ml	5 tablets	2 1/2 tablets	1 tablet
72 - 95	33 - 43	480 mg		3 teaspoons 15 ml	6 tablets	3 tablets	1 1/2 tablets
96 +	44 +	650 mg		4 teaspoons 20 ml	8 tablets	4 tablets	2 tablets

### Advil®/ Motrin® (Ibuprofen) dosing chart for home use

· Advil®/Motrin® (Ibuprofen) may be given every 6 hours.

· Advil®/ Motrin® (Ibuprofen) is not to be given to infants under 6 months of age.

Weight in Pounds (lb)	Weight in Kilograms (kg)	Dosage in Mg	Infant Drops 50 mg/1.25 ml 40 mg/ml	Children's Liquid 100 mg/5 ml 20 mg/ml	Chewable Tablets 50 mg tabs	Junior Strength Tablets 100 mg tabs	Adult Strength Tablets 200 mg tabs
12 - 17	5 - 7	50 mg	1.25 ml	1/2 teaspoon 2.5 ml			
18 - 23	8 - 10	75 mg	1.875 ml	3/4 teaspoon 3.75 ml			
24 - 35	11 - 15	100 mg	2.5 ml	1 teaspoon 5 ml	2 tablets	1 tablet	
36 - 47	16 - 21	150 mg	3.75 ml	1 1/2 teaspoons 7.5 ml	3 tablets	1 1/2 tablets	
48 - 59	22 - 26	200 mg	5 ml	2 teaspoons 10 ml	4 tablets	2 tablets	1 tablet
60 - 71	27 - 32	250 mg		2 1/2 teaspoons 12.5 ml	5 tablets	2 1/2 tablets	1 tablet
72 - 95	33 - 43	300 mg		3 teaspoons 15 ml	6 tablets	3 tablets	1 1/2 tablets
96 +	44 +	400 mg		4 teaspoons 20 ml	8 tablets	4 tablets	2 tablets

· Although generally not recommended, if your health care provider instructs you to alternate Acetaminophen and Ibuprofen, then alternate doses every four hours. Alternate medications only for 24 hours or less then return to using a single product.

· Children under 18 years old should never take aspirin or aspirin containing medicines due to the risk of Reye's Syndrome, a severe encephalitis like illness.



## Medicamentos para aliviar la fiebre/dolor para bebés/niños

\* Compruebe la concentración y el tipo de medicamento en el frasco que está utilizando antes de dosificar a su hijo \*

### Tabla de dosificación de Tylenol® (Acetaminofén) para uso doméstico

· Tylenol® (Acetaminofén) se puede administrar a cada 4 horas

· Para un niño con vómitos o convulsiones que no puede tomar paracetamol oral hay supositorios rectales disponibles en dosis 120 mg y (Utilice la tabla de dosificación a continuación para encontrar la dosis en mg necesaria)

Peso en libras	Peso en kilogramos	Dosis en mg	Gotas para bebés 160 mg/5 ml	Gotas para Niños 160 mg/5 ml	Tabletas masticables 80 mg tabs	Tabletas para adolescentes 160 mg tabs	Tabletas para adultos 325 mg tabs
6 - 11	3 - 5	40 mg	1.25 ml	1/4 teaspoon 1.25 ml			
12 - 17	6 - 7	80 mg	2.5 ml	1/2 teaspoon 2.5 ml			
18 - 23	8 - 10	120 mg	3.75 ml	3/4 teaspoon 3.75 ml			
24 - 35	11 - 15	160 mg	5 ml	1 teaspoon 5 ml	2 tablets	1 tablet	
36 - 47	16 - 21	240 mg		1 1/2 teaspoons 7.5 ml	3 tablets	1 1/2 tablets	
48 - 59	22 - 26	320 mg		2 teaspoons 10 ml	4 tablets	2 tablets	1 tablet
60 - 71	27 - 32	400 mg		2 1/2 teaspoons 12.5 ml	5 tablets	2 1/2 tablets	1 tablet
72 - 95	33 - 43	480 mg		3 teaspoons 15 ml	6 tablets	3 tablets	1 1/2 tablets
96 +	44 +	650 mg		4 teaspoons 20 ml	8 tablets	4 tablets	2 tablets

### Tabla de dosificación de Advil®/ Motrin® (Ibuprofeno) para uso doméstico

· Advil/Motrin (Ibuprofeno) se puede administrar a cada 6 horas

· No debe administrarse a bebés menores de 6 meses

Peso en libras (lb)	Peso en kilogramos (kg)	Dosis en mg	Gotas para Bebés 50 mg/1.25 ml	Gotas para Niños 100 mg/5 ml	Tabletas masticables 50 mg tabs	Tabletas para adolescentes 100 mg tabs	Tabletas para adultos 200 mg tabs
12 - 17	5 - 7	50 mg	1.25 ml	1/2 teaspoon 2.5 ml			
18 - 23	8 - 10	75 mg	1.875 ml	3/4 teaspoon 3.75 ml			
24 - 35	11 - 15	100 mg	2.5 ml	1 teaspoon 5 ml	2 tablets	1 tablet	
36 - 47	16 - 21	150 mg	3.75 ml	1 1/2 teaspoons 7.5 ml	3 tablets	1 1/2 tablets	
48 - 59	22 - 26	200 mg	5 ml	2 teaspoons 10 ml	4 tablets	2 tablets	1 tablet
60 - 71	27 - 32	250 mg		2 1/2 teaspoons 12.5 ml	5 tablets	2 1/2 tablets	1 tablet
72 - 95	33 - 43	300 mg		3 teaspoons 15 ml	6 tablets	3 tablets	1 1/2 tablets
96 +	44 +	400 mg		4 teaspoons 20 ml	8 tablets	4 tablets	2 tablets

· Aunque generalmente no se recomienda, pero si su proveedor de atención médica le indica el paracetamol y el ibuprofeno, entonces alterne la dosis a cada 4 horas. Los medicamentos alternativos solo deben administrarse durante 24 horas, después regresa a un solo producto.

· Los niños menores de 18 años nunca deben tomar aspirina o medicina que contengan medicamentos debido al riesgo del síndrome de Reye, que es una encefalitis grave como la enfermedad.