



VACCINE SCHEDULE

NURSERY	Hepatitis B #1
ONE MONTH	<i>catch-up any missing immunizations</i>
TWO MONTHS	Pentacel #1 (DTaP-IPV/HIB), Prevnar #1, Rotateq #1
THREE MONTHS	Hepatitis B #2 ¹
FOUR MONTHS	Pentacel #2 (DTaP-IPV/HIB), Prevnar #2, Rotateq #2
SIX MONTHS	Pentacel #3 (DTaP-IPV/HIB), Prevnar #3, Rotateq #3
NINE MONTHS	Hepatitis B #3
ONE YEAR (must be 1 year)	MMR, Varivax, Hepatitis A #1
FIFTEEN MONTHS	Pentacel #4 (DTaP-IPV/HIB), Prevnar #4
EIGHTEEN MONTHS	<i>Hepatitis A #2 if appropriate ²</i>
TWO YEARS	<i>Hepatitis A #2 if not given at 18 months</i>
FOUR-SIX YEARS	DTaP-IPV, MMR/V
NINE YEARS	HPV 9 #1 (Gardasil) ³
ELEVEN-TWELVE YEARS	Tdap ⁴, Meningococcal ACYW #1
SIXTEEN YEARS	Meningococcal ACYW #1
SIXTEEN-EIGHTEEN YEARS	Meningococcal B #1/#2 (Bexsero) ⁵

Notes:

1. If an infant is born to a mother with Hepatitis B, the 0-1-6 month schedule should be followed
2. Hepatitis A vaccine may be administered once the patient is over 12 months of age; dose #1 and #2 are given 6-18 months apart
3. HPV vaccine series can be started at age 9 years
 - if the first dose is given before the patient is 15 years old, only one other dose is needed; dose #1 and #2 should be separated by at least 6 months
 - if a patient is 15 years or older when the first dose is given, 3 total doses will be required to complete the series
4. If a child is 10 years old and tetanus prophylaxis is needed, Tdap should be given
5. Meningococcal B #1 and #2 are given at least 6 months apart

Additional notes:

Flu vaccine is given seasonally, from August through April (depending on supply); patients who receive their first dose before the age of 9 years are recommended to receive 2 doses in the same season, at least 4 weeks apart; if 2 doses cannot be given in the same season, it should be done the following season

RSV immunization is administered to infants 0-8 months of age in October through March if Mom did not receive RSV maternal vaccine or if the baby is born less than 2 weeks after Mom received the vaccine; Mom is recommended to receive the maternal RSV vaccine at 32-36 weeks of gestation

Adolescents 13 years and older, who have not received any or all of the vaccines designated for the 9-12 year age group, should be given these vaccines; those who have not yet started the Hepatitis B series should receive this vaccine also

Yearly Well Visits from the ages of 3 – 21 years are expected in order to be a patient in good standing

Those patients who do not have yearly Well Visits will not be eligible to have

- *forms completed and/or signed off by our office*
- *medication refills*
- *administration of certain medications and vaccinations in office*
- *mental health therapy or psychiatric services in office*