Coronavirus (COVID-19) Self-Monitoring Information

This form may be utilized if you have been exposed to an under assumed or confirmed case to an individual with COVID-19.

Guidance for persons who choose to self-monitor:

This guidance is to help you monitor your health for up to 14 days because you may have been exposed to coronavirus. This does NOT mean that you will get sick with coronavirus. You may choose to monitor your health so that if you develop symptoms, you can seek medical care and reduce the spread of illness to others. Based on what is known from other coronavirus infections, 14 days is likely the longest time between when you were last exposed to coronavirus and when symptoms begin.

What are the signs and symptoms of coronavirus?

Per the CDC, these symptoms may appear 2-14 days after exposure (Fever, cough, shortness of breath). Some people also report more rarely sore throat, muscle aches, and abdominal discomfort. It is important to note that these symptoms are associated with many other illnesses.

How should you monitor your health during this time period?

Use the designated page of this form to record your temperature and possible symptoms twice every day.

Practicing protective and careful measures during this time period.

- For more information, <u>https://www.cdc.gov/coronavirus/2019-ncov/downloads/sick-with-2019-nCoV-fact-sheet.pdf</u>
- Please see the following page regarding CDC's recommendation
- Recent recommendations are to wear a cloth face covering over your nose and mouth, if sick

Instructions for monitoring your temperature and symptoms

- 1. Take your temperature orally (by mouth) with a thermometer twice a day in the morning and evening.
- 2. Write down your temperature on the form with the corresponding time
- 3. Checkmark any of the symptoms of the novel coronavirus or specify any other symptoms you may be experiencing

What should I do if I develop symptoms/ become ill during this monitoring period?

Call ahead. Call your primary care provider or one of the numbers below before you seek care. If your symptoms are mild, it is possible that you may be able to remain at home. If it is determined that you may need medical attention, they will provide information regarding how to seek care in the safest way possible.

If you feel your symptoms are at EMERGENCY LEVELS i.e. shortness of breath, fever over 102 F, chest pain, etc., that cannot wait until the morning, call 911 and explain that you are being monitored for potential exposure to the coronavirus and need emergency medical care.

Name of Staff: Date of Exposure: Exposure Category: Low/ Medium/ High							Department of Health and Human Services (DHHS) – 800-860-0620					
Date:	Temperature (AM/ PM)		Coughing Checkmark(✓)	Sore Throat Checkmark(✓)	Chills Checkmark(✓)		Abdominal Discomfort Checkmark(✓)	Diarrhea Checkmark(✓	Other Shortness of Breath Checkmark(✓)	Muscle Aches Checkmark(✓)	Other: Specify	
Day 1:	Time AM	Time Time AM PM										
Day 2:	Time	Time PM										
Day 3:	Time AM	Time PM										
Day 4:	Time AM	Time PM										
Day 5:	Time AM	Time PM										
Day 6:	Time AM	Time PM										
Day 7:	Time AM	Time PM										
Day 8:	Time AM	Time PM										
Day 9:	Time AM	Time PM										
Day 10:	Time AM	Time PM										
Day 11:	Time AM	Time PM										
Day 12:	Time AM	Time PM										
Day 13:	Time AM	Time PM										
Day 14:	Time AM	Time PM										

Steps to help prevent the spread of COVID-19 if you are sick

FOLLOW THE STEPS BELOW: If you are sick with COVID-19 or think you might have COVID-19, follow the steps below to help protect other people in your home and community.

Stay home except to get medical care

- **Stay home:** Most people with COVID-19 have mild illness and are able to recover at home without medical care. Do not leave your home, except to get medical care. Do not visit public areas.
- **Stay in touch with your doctor.** Call before you get medical care. Be sure to get care if you have trouble breathing, or have any other emergency warning signs, or if you think it is an emergency.
- Avoid public transportation: Avoid using public transportation, ride-sharing, or taxis.

Separate yourself from other people in your home, this is known as home isolation



- **Stay away from others:** As much as possible, stay away from others. You should stay in a specific "sick room" if possible, and away from other people in your home. Use a separate bathroom, if available.
 - See COVID-19 and Animals is you have questions about pets. <u>https://www.cdc.gov/coronavirus/2019-ncov/faq.</u> <u>html#COVID19animals</u>

Call ahead before visiting your doctor

• **Call ahead:** Many medical visits for routine care are being postponed or done by phone or telemedicine.



• If you have a medical appointment that cannot be postponed, call your doctor's office, and tell them you have or may have COVID-19. This will help the office protect themselves and other patients.

If you are sick wear a facemask in the following situations, if available.

If you are sick: You should wear a facemask, if available, when you are around other people (including before you enter a healthcare provider's office).



• **If you are caring for others:** If the person who is sick is not able to wear a facemask (for example, because it causes trouble breathing), then as their caregiver, you should wear a facemask when in the same room with them. Visitors, other than caregivers, are not recommended.

Note: During a public health emergency, facemasks may be reserved for healthcare workers. You may need to improvise a facemask using a scarf or bandana.

Cover your coughs and sneezes

• **Cover:** Cover your mouth and nose with a tissue when you cough or sneeze.



- **Dispose:** Throw used tissues in a lined trash can.
- Wash hands: Immediately wash your hands with soap and water for at least 20 seconds. If soap and water are not available, clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol.

Clean your hands often

• Wash hands: Wash your hands often with soap and water for at least 20 seconds. This is especially important after blowing your nose, coughing, or sneezing; going to the bathroom; and before eating or preparing food.



- Hand sanitizer: If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry.
- **Soap and water:** Soap and water are the best option, especially if hands are visibly dirty.
- **Avoid touching:** Avoid touching your eyes, nose, and mouth with unwashed hands.

Avoid sharing personal household items

• **Do not share:** Do not share dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people in your home.



• Wash thoroughly after use: After using these items, wash them thoroughly with soap and water or put in the dishwasher.



cdc.gov/coronavirus

CS 316120-A 04/02/2020

Clean all "high-touch" surfaces everyday

Clean high-touch surfaces in your isolation area ("sick room" and bathroom) every day; let a caregiver clean and disinfect high-touch surfaces in other areas of the home.



- **Clean and disinfect:** Routinely clean high-touch surfaces in your "sick room" and bathroom. Let someone else clean and disinfect surfaces in common areas, but not your bedroom and bathroom.
 - If a caregiver or other person needs to clean and disinfect a sick person's bedroom or bathroom, they should do so on an as-needed basis. The caregiver/other person should wear a mask and wait as long as possible after the sick person has used the bathroom.
- High-touch surfaces include phones, remote controls, counters, tabletops, doorknobs, bathroom fixtures, toilets, keyboards, tablets, and bedside tables.
- Clean and disinfect areas that may have blood, stool, or body fluids
 on them.
- Household cleaners and disinfectants: Clean the area or item with soap and water or another detergent if it is dirty. Then, use a household disinfectant.
 - Be sure to follow the instructions on the label to ensure safe and effective use of the product. Many products recommend keeping the surface wet for several minutes to ensure germs are killed. Many also recommend precautions such as wearing gloves and making sure you have good ventilation during use of the product.
 - Most EPA-registered household disinfectants should be effective. A full list of disinfectants can be found hereexternal icon.

Monitor your symptoms

• Common symptoms of COVID-19 include fever and cough. Trouble breathing is a more serious symptom that means you should get medical attention.



- If you are having trouble breathing, seek medical attention, but call first.
 - Call your doctor or emergency room before going in and tell them your symptoms. They will tell you what to do.
- Wear a facemask: If available, put on a facemask before you enter the building. If you can't put on a facemask, cover your coughs and sneezes. Try to stay at least 6 feet away from other people. This will help protect the people in the office or waiting room.
- Follow care instructions from your healthcare provider and local health department: Your local health authorities will give instructions on checking your symptoms and reporting information.

If you develop **emergency warning signs** for COVID-19 get **medical attention immediately.**

Emergency warning signs include*:

- Trouble breathing
- Persistent pain or pressure in the chest
- New confusion or inability to arouse
- Bluish lips or face

*This list is not all inclusive. Please consult your medical provider for any other symptoms that are severe or concerning.

Call 911 if you have a medical emergency: If you have a medical emergency and need to call 911, notify the operator that you have or think you might have, COVID-19. If possible, put on a facemask before medical help arrives.

How to discontinue home isolation

 People with COVID-19 who have stayed home (home isolated) can stop home isolation under the following conditions:



- If you will not have a test to determine if you are still contagious, you can leave home after these three things have happened:
 - You have had no fever for at least 72 hours (that is three full days of no fever without the use medicine that reduces fevers)
 AND
 - other symptoms have improved (for example, when your cough or shortness of breath have improved) AND
 - at least 7 days have passed since your symptoms first appeared
- If you will be tested to determine if you are still contagious, you can leave home after these three things have happened:
 - You no longer have a fever (without the use medicine that reduces fevers) AND
 - other symptoms have improved (for example, when your cough or shortness of breath have improved) AND
 - you received two negative tests in a row, 24 hours apart. Your doctor will follow CDC guidelines.

In all cases, follow the guidance of your healthcare provider and local health department. The decision to stop home isolation should be made in consultation with your healthcare provider and state and local health departments. Local decisions depend on local circumstances.

More information is available <u>https://www.cdc.gov/</u> coronavirus/2019-ncov/hcp/disposition-in-home-patients.html.

Additional information for healthcare providers: <u>Interim</u> <u>Healthcare Infection Prevention and Control Recommendations</u> <u>for Persons Under Investigation for 2019 Novel Coronavirus.</u>