Generalized Anxiety Disorder Screener (GAD-7)

Date:

	er the last 2 weeks, how often have you been	Not at all	Several	More than	Nearly
bot	thered by the following problems?		Days	half the days	every day
1.	Feeling nervous, anxious or on edge	0	1	2	3
2.	Not being able to stop or control worrying	0	1	2	3
3.	Worrying too much about different things	0	1	2	3
4.	Trouble relaxing	0	1	2	3
5.	Being so restless that it is hard to sit still	0	1	2	3
6.	Becoming easily annoyed or irritated	0	1	2	3
7.	Feeling afraid as if something awful might happen	0	1	2	3
		Add columns			
		Total Score			
8.	If you checked off any problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?	Not difficult at all	Somewhat difficult	Very difficult	Extremely difficult

When did the symptoms begin? _			
Patient Name:			
DOB:			

Generalized Anxiety Disorder Screener (GAD-7)

Scoring and Interpretation:

GAD-2 Score*	Provisional Diagnosis	
0-2	None	
3-6	Probable anxiety disorder	
GAD-7 Score	Provisional Diagnosis	
0-7	None	
8+	Probable anxiety disorder	

^{*}GAD-2 is the first 2 questions of the GAD-7

References:

- Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. Archives of internal medicine. May 22 2006;166(10):1092-1097. PMID: 16717171
- Kroenke K, Spitzer RL, Williams JB, Monahan PO, Lowe B. Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. Annals of internal medicine. Mar 6 2007;146(5):317-325. PMID: 17339617
- Lowe B, Decker O, Muller S, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. Medical care. Mar 2008;46(3):266-274. PMID: 18388841