

**WARNING: This product contains nicotine.
Nicotine is an addictive chemical.**

VAPOR[™]
EFX

HOW TO USE

Micelle Nicotine, NanoMist Spray

Sublingual Aerosol



1. Holding the VaporEFX[™] device upright, turn the **cap clockwise** on the base to unlock the **child-resistant cap**. You will notice two faint arrows on the cap and base that will align, letting you know that the cap is in the ready to remove position. **Remove the cap.**
2. Bring the Vapor device to your mouth; with your mouth open, **point the black actuator nozzle to the inside of your mouth** and press firmly down on the spray actuator to release the NanoMist[™] spray directly under your tongue, or to the inside of your cheeks. Be certain to hold the VaporEFX device close to your mouth to avoid direct contact with your eyes or lips.
3. Repeat this process based on your desired amount of nicotine. Please note that each VaporEFX spray delivers a precise, metered amount of micelle nicotine (.5 mg or 1.0 mg). **Repeat as desired.**
4. For best results, and greater nicotine absorption, **wait approximately one-minute** before swallowing. Most users find that the NanoMist spray is quickly absorbed into the mucosal lining of your mouth.
5. Users may notice a cloud like Vapor when spraying the NanoMist into their mouth. **This is a normal, thermal reaction** that takes place when your body temperature is warmer than the colder temperature of the NanoMist spray propellant.
6. To secure and lock the cap, place the cap back onto the base with the arrows aligned. **Turn the cap counterclockwise**, less than a half of a turn, until you hear or feel a click. **The cap is now secured and locked.**



For Additional Product Information, Please Contact:

E: Info@Air2Vapor.com

O: (303) 736-2874

Air 2, LLC 8100 S. Akron St. Suite 316, Centennial, Colorado USA