

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	91.6%
	Follow-up at Six Months	Expected	54.4%
	Response at Six Months	Expected	24.9%
	Remission at Six Months	Expected	12.4%
	Follow-up at 12 Months	Above Expected	60.7%
	Response at 12 Months	Expected	27.2%
	Remission at 12 Months	Expected	15.7%
Adolescent Mental Health and/or Depression Screening*		Below	81.7%
Colorectal Cancer Screening		Above Expected	78.6%
Optimal Asthma Control - Adults		Above Expected	74.3%
Optimal Asthma Control - Children		Above Expected	66.9%
Optimal Diabetes Care		Above Expected	56.5%
HbA1c Control		Above Expected	81.7%
BP Control		Expected	81.6%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	91.0%
Tobacco-Free		Above Expected	91.1%
Optimal Vascular Care		Expected	58.3%
BP Control		Expected	83.3%
Daily Aspirin Use		Below Expected	81.8%
Statin Use		Expected	92.9%
Tobacco-Free		Above Expected	90.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).