

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Below	47.2%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	38.2%
	Follow-up at Six Months	Below Expected	0.0%
	Response at Six Months	Below Expected	0.0%
	Remission at Six Months	Expected	0.0%
	Follow-up at 12 Months	Below Expected	2.9%
	Response at 12 Months	Below Expected	0.0%
	Remission at 12 Months	Expected	0.0%
Adolescent Mental Health and/or Depression Screening*		Below	80.9%
Colorectal Cancer Screening		Above Expected	74.7%
Optimal Asthma Control - Adults		Below Expected	0.0%
Optimal Asthma Control - Children		Below Expected	0.0%
Optimal Diabetes Care		Above Expected	51.0%
HbA1c Control		Above Expected	77.2%
BP Control		Expected	80.5%
Daily Aspirin Use		Expected	98.6%
Statin Use		Expected	89.4%
Tobacco-Free		Above Expected	91.4%
Optimal Vascular Care		Expected	55.5%
BP Control		Expected	80.8%
Daily Aspirin Use		Expected	86.4%
Statin Use		Below Expected	85.4%
Tobacco-Free		Above Expected	87.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).