

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	39.6%
	Follow-up at Six Months	Below Expected	18.0%
	Response at Six Months	Below Expected	3.9%
	Remission at Six Months	Below Expected	0.0%
	Follow-up at 12 Months	Below Expected	18.0%
	Response at 12 Months	Below Expected	1.3%
	Remission at 12 Months	Below Expected	0.0%
Adolescent Mental Health and/or Depression Screening*		Below	56.3%
Colorectal Cancer Screening		Below Expected	31.3%
Optimal Asthma Control - Adults		Below Expected	9.4%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	18.3%
HbA1c Control		Below Expected	58.1%
BP Control		Below Expected	59.1%
Daily Aspirin Use		Expected	99.5%
Statin Use		Expected	86.6%
Tobacco-Free		Below Expected	62.4%
Optimal Vascular Care		Below Expected	22.0%
BP Control		Below Expected	50.9%
Daily Aspirin Use		Expected	89.8%
Statin Use		Expected	93.2%
Tobacco-Free		Below Expected	55.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).