

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	67.6%
	Follow-up at Six Months	Below Expected	35.7%
	Response at Six Months	Expected	15.5%
	Remission at Six Months	Expected	13.1%
	Follow-up at 12 Months	Below Expected	35.7%
	Response at 12 Months	Expected	15.5%
	Remission at 12 Months	Expected	10.7%
Adolescent Mental Health and/or Depression Screening*		Below	82.5%
Colorectal Cancer Screening		Below Expected	65.1%
Optimal Asthma Control - Adults		Expected	40.4%
Optimal Asthma Control - Children		Expected	40.0%
Optimal Diabetes Care		Expected	41.9%
HbA1c Control		Expected	77.4%
BP Control		Expected	76.7%
Daily Aspirin Use		Expected	99.6%
Statin Use		Below Expected	81.0%
Tobacco-Free		Expected	83.9%
Optimal Vascular Care		Expected	65.9%
BP Control		Expected	83.7%
Daily Aspirin Use		Expected	93.0%
Statin Use		Expected	89.9%
Tobacco-Free		Expected	85.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).