

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	87.0%
	Follow-up at Six Months	Expected	59.6%
	Response at Six Months	Expected	21.3%
	Remission at Six Months	Expected	12.1%
	Follow-up at 12 Months	Expected	58.0%
	Response at 12 Months	Expected	24.2%
	Remission at 12 Months	Expected	14.3%
Adolescent Mental Health and/or Depression Screening*		Below	88.3%
Colorectal Cancer Screening		Expected	74.1%
Optimal Asthma Control - Adults		Above Expected	69.3%
Optimal Asthma Control - Children		Expected	61.3%
Optimal Diabetes Care		Above Expected	54.2%
HbA1c Control		Expected	75.8%
BP Control		Expected	81.8%
Daily Aspirin Use		Above Expected	99.6%
Statin Use		Expected	87.6%
Tobacco-Free		Above Expected	92.0%
Optimal Vascular Care		Above Expected	65.2%
BP Control		Expected	84.8%
Daily Aspirin Use		Expected	87.7%
Statin Use		Expected	93.6%
Tobacco-Free		Expected	89.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).