

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	68.8%
	Follow-up at Six Months	Expected	54.7%
	Response at Six Months	Expected	11.3%
	Remission at Six Months	Expected	11.3%
	Follow-up at 12 Months	Expected	47.2%
	Response at 12 Months	Expected	11.3%
	Remission at 12 Months	Below Expected	1.9%
Adolescent Mental Health and/or Depression Screening*		Average	90.9%
Colorectal Cancer Screening		Expected	70.8%
Optimal Asthma Control - Adults		Below Expected	16.7%
Optimal Asthma Control - Children		Below Expected	11.8%
Optimal Diabetes Care		Expected	43.6%
HbA1c Control		Expected	77.0%
BP Control		Expected	78.4%
Daily Aspirin Use		Above Expected	100.0%
Statin Use		Expected	87.1%
Tobacco-Free		Below Expected	75.6%
Optimal Vascular Care		Below Expected	41.2%
BP Control		Expected	75.4%
Daily Aspirin Use		Expected	89.8%
Statin Use		Expected	94.7%
Tobacco-Free		Below Expected	62.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).