

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	69.8%
	Follow-up at Six Months	Below Expected	26.5%
	Response at Six Months	Below Expected	9.7%
	Remission at Six Months	Below Expected	4.1%
	Follow-up at 12 Months	Below Expected	31.1%
	Response at 12 Months	Expected	13.3%
	Remission at 12 Months	Expected	6.1%
Adolescent Mental Health and/or Depression Screening*		Below	82.3%
Colorectal Cancer Screening		Expected	70.0%
Optimal Asthma Control - Adults		Below Expected	17.0%
Optimal Asthma Control - Children		Expected	33.3%
Optimal Diabetes Care		Expected	45.6%
HbA1c Control		Expected	71.2%
BP Control		Expected	85.0%
Daily Aspirin Use		Expected	98.1%
Statin Use		Expected	89.1%
Tobacco-Free		Expected	85.0%
Optimal Vascular Care		Expected	57.2%
BP Control		Expected	82.2%
Daily Aspirin Use		Expected	85.5%
Statin Use		Expected	94.6%
Tobacco-Free		Expected	83.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).