

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	92.5%
	Follow-up at Six Months	Expected	42.9%
	Response at Six Months	Expected	16.1%
	Remission at Six Months	Expected	10.7%
	Follow-up at 12 Months	Expected	32.1%
	Response at 12 Months	Expected	10.7%
	Remission at 12 Months	Expected	7.1%
Adolescent Mental Health and/or Depression Screening*		Average	93.8%
Colorectal Cancer Screening		Below Expected	44.2%
Optimal Asthma Control - Adults		Below Expected	12.1%
Optimal Asthma Control - Children		Below Expected	16.4%
Optimal Diabetes Care		Expected	39.9%
HbA1c Control		Below Expected	55.5%
BP Control		Expected	84.6%
Daily Aspirin Use		Expected	99.5%
Statin Use		Above Expected	92.3%
Tobacco-Free		Above Expected	92.0%
Optimal Vascular Care		Expected	40.6%
BP Control		Expected	68.8%
Daily Aspirin Use		Expected	75.0%
Statin Use		Expected	90.6%
Tobacco-Free		Expected	81.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).