

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	58.8%
	Follow-up at Six Months	Expected	31.4%
	Response at Six Months	Expected	11.4%
	Remission at Six Months	Expected	7.1%
	Follow-up at 12 Months	Expected	34.3%
	Response at 12 Months	Expected	15.7%
	Remission at 12 Months	Expected	8.6%
Adolescent Mental Health and/or Depression Screening*		Below	73.2%
Colorectal Cancer Screening		Above Expected	76.9%
Optimal Asthma Control - Adults		Below Expected	24.8%
Optimal Asthma Control - Children		Expected	28.6%
Optimal Diabetes Care		Below Expected	38.1%
HbA1c Control		Expected	73.2%
BP Control		Expected	78.7%
Daily Aspirin Use		Below Expected	95.9%
Statin Use		Expected	83.8%
Tobacco-Free		Expected	85.0%
Optimal Vascular Care		Expected	47.1%
BP Control		Expected	84.1%
Daily Aspirin Use		Below Expected	73.5%
Statin Use		Expected	89.9%
Tobacco-Free		Expected	84.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).