

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	83.5%
	Follow-up at Six Months	Below Expected	30.2%
	Response at Six Months	Below Expected	9.6%
	Remission at Six Months	Below Expected	4.8%
	Follow-up at 12 Months	Below Expected	35.3%
	Response at 12 Months	Below Expected	12.2%
	Remission at 12 Months	Expected	7.4%
Adolescent Mental Health and/or Depression Screening*		Below	80.6%
Colorectal Cancer Screening		Above Expected	73.2%
Optimal Asthma Control - Adults		Below Expected	43.2%
Optimal Asthma Control - Children		Expected	39.0%
Optimal Diabetes Care		Below Expected	39.3%
HbA1c Control		Below Expected	67.3%
BP Control		Above Expected	84.8%
Daily Aspirin Use		Below Expected	94.3%
Statin Use		Below Expected	81.2%
Tobacco-Free		Expected	85.4%
Optimal Vascular Care		Below Expected	30.8%
BP Control		Expected	83.2%
Daily Aspirin Use		Below Expected	53.9%
Statin Use		Below Expected	79.1%
Tobacco-Free		Expected	78.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).