

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	45.1%
	Response at Six Months	Expected	20.8%
	Remission at Six Months	Expected	14.6%
	Follow-up at 12 Months	Expected	46.5%
	Response at 12 Months	Expected	20.1%
	Remission at 12 Months	Expected	11.1%
Adolescent Mental Health and/or Depression Screening*		Below	85.0%
Colorectal Cancer Screening		Above Expected	75.5%
Optimal Asthma Control - Adults		Expected	47.9%
Optimal Asthma Control - Children		Expected	40.0%
Optimal Diabetes Care		Expected	52.4%
HbA1c Control		Expected	76.9%
BP Control		Above Expected	87.2%
Daily Aspirin Use		Expected	99.1%
Statin Use		Expected	86.7%
Tobacco-Free		Expected	88.6%
Optimal Vascular Care		Expected	57.3%
BP Control		Expected	83.7%
Daily Aspirin Use		Expected	84.6%
Statin Use		Expected	87.2%
Tobacco-Free		Expected	85.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).