

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	32.5%
	Follow-up at Six Months	Below Expected	11.9%
	Response at Six Months	Below Expected	3.4%
	Remission at Six Months	Below Expected	0.9%
	Follow-up at 12 Months	Below Expected	29.7%
	Response at 12 Months	Expected	11.0%
	Remission at 12 Months	Below Expected	3.4%
Adolescent Mental Health and/or Depression Screening*		Below	86.5%
Colorectal Cancer Screening		Below Expected	66.4%
Optimal Asthma Control - Adults		Below Expected	0.0%
Optimal Asthma Control - Children		Below Expected	0.0%
Optimal Diabetes Care		Below Expected	32.2%
HbA1c Control		Expected	78.0%
BP Control		Below Expected	67.4%
Daily Aspirin Use		Expected	97.6%
Statin Use		Below Expected	68.8%
Tobacco-Free		Below Expected	77.4%
Optimal Vascular Care		Below Expected	38.8%
BP Control		Below Expected	67.0%
Daily Aspirin Use		Below Expected	80.9%
Statin Use		Below Expected	84.1%
Tobacco-Free		Below Expected	74.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).