

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	55.2%
	Response at Six Months	Expected	22.4%
	Remission at Six Months	Expected	12.6%
	Follow-up at 12 Months	Expected	44.8%
	Response at 12 Months	Expected	13.3%
	Remission at 12 Months	Expected	4.9%
Adolescent Mental Health and/or Depression Screening*		Below	53.8%
Colorectal Cancer Screening		Above Expected	73.6%
Optimal Asthma Control - Adults		Expected	60.3%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	62.5%
HbA1c Control		Expected	77.2%
BP Control		Above Expected	95.1%
Daily Aspirin Use		Expected	99.4%
Statin Use		Expected	91.1%
Tobacco-Free		Above Expected	88.9%
Optimal Vascular Care		Above Expected	68.7%
BP Control		Above Expected	95.3%
Daily Aspirin Use		Expected	87.4%
Statin Use		Expected	93.9%
Tobacco-Free		Expected	83.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).