

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	52.4%
	Response at Six Months	Expected	15.2%
	Remission at Six Months	Expected	7.8%
	Follow-up at 12 Months	Expected	45.5%
	Response at 12 Months	Below Expected	11.7%
	Remission at 12 Months	Below Expected	4.3%
Adolescent Mental Health and/or Depression Screening*		Below	89.1%
Colorectal Cancer Screening		Expected	64.9%
Optimal Asthma Control - Adults		Below Expected	26.2%
Optimal Asthma Control - Children		Expected	38.5%
Optimal Diabetes Care		Expected	45.4%
HbA1c Control		Expected	67.3%
BP Control		Above Expected	84.2%
Daily Aspirin Use		Expected	99.2%
Statin Use		Expected	90.9%
Tobacco-Free		Above Expected	85.1%
Optimal Vascular Care		Expected	56.3%
BP Control		Expected	84.4%
Daily Aspirin Use		Expected	87.9%
Statin Use		Expected	92.9%
Tobacco-Free		Expected	79.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).