

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	90.0%
	Follow-up at Six Months	Expected	53.6%
	Response at Six Months	Expected	21.1%
	Remission at Six Months	Expected	9.7%
	Follow-up at 12 Months	Above Expected	61.9%
	Response at 12 Months	Above Expected	23.9%
	Remission at 12 Months	Above Expected	15.3%
Adolescent Mental Health and/or Depression Screening*		Below	89.4%
Colorectal Cancer Screening		Expected	67.7%
Optimal Asthma Control - Adults		Above Expected	59.5%
Optimal Asthma Control - Children		Above Expected	56.9%
Optimal Diabetes Care		Expected	43.2%
HbA1c Control		Expected	71.8%
BP Control		Expected	78.6%
Daily Aspirin Use		Above Expected	99.6%
Statin Use		Expected	87.5%
Tobacco-Free		Expected	83.4%
Optimal Vascular Care		Expected	55.0%
BP Control		Expected	81.2%
Daily Aspirin Use		Above Expected	90.2%
Statin Use		Expected	93.2%
Tobacco-Free		Expected	78.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).