

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	91.8%
	Follow-up at Six Months	Above Expected	81.7%
	Response at Six Months	Above Expected	37.3%
	Remission at Six Months	Above Expected	23.9%
	Follow-up at 12 Months	Above Expected	75.4%
	Response at 12 Months	Above Expected	36.6%
	Remission at 12 Months	Above Expected	25.4%
Adolescent Mental Health and/or Depression Screening*		Average	94.7%
Colorectal Cancer Screening		Below Expected	67.9%
Optimal Asthma Control - Adults		Above Expected	71.9%
Optimal Asthma Control - Children		Above Expected	72.8%
Optimal Diabetes Care		Above Expected	58.0%
HbA1c Control		Expected	77.3%
BP Control		Expected	83.5%
Daily Aspirin Use		Above Expected	99.8%
Statin Use		Above Expected	93.9%
Tobacco-Free		Expected	88.6%
Optimal Vascular Care		Expected	64.1%
BP Control		Above Expected	90.2%
Daily Aspirin Use		Expected	87.0%
Statin Use		Expected	94.3%
Tobacco-Free		Expected	84.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).