

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	46.9%
	Follow-up at Six Months	Below Expected	23.3%
	Response at Six Months	Expected	10.0%
	Remission at Six Months	Expected	3.3%
	Follow-up at 12 Months	Expected	28.3%
	Response at 12 Months	Expected	10.0%
	Remission at 12 Months	Expected	5.0%
Adolescent Mental Health and/or Depression Screening*		Below	81.6%
Colorectal Cancer Screening		Expected	64.0%
Optimal Asthma Control - Adults		Below Expected	16.6%
Optimal Asthma Control - Children		Below Expected	16.7%
Optimal Diabetes Care		Below Expected	36.3%
HbA1c Control		Expected	64.8%
BP Control		Expected	83.0%
Daily Aspirin Use		Below Expected	94.9%
Statin Use		Below Expected	77.9%
Tobacco-Free		Expected	84.3%
Optimal Vascular Care		Below Expected	31.5%
BP Control		Expected	84.5%
Daily Aspirin Use		Below Expected	62.0%
Statin Use		Below Expected	80.3%
Tobacco-Free		Expected	75.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).