

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	81.8%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	94.4%
Colorectal Cancer Screening		Above Expected	80.8%
Optimal Asthma Control - Adults		Expected	45.5%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	48.6%
HbA1c Control		Above Expected	79.1%
BP Control		Above Expected	89.5%
Daily Aspirin Use		Below Expected	92.9%
Statin Use		Expected	88.6%
Tobacco-Free		Expected	78.1%
Optimal Vascular Care		Expected	47.7%
BP Control		Above Expected	92.0%
Daily Aspirin Use		Below Expected	70.5%
Statin Use		Expected	93.3%
Tobacco-Free		Expected	72.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).