

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	79.4%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Expected	77.5%
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	50.7%
HbA1c Control		Expected	82.9%
BP Control		Expected	80.8%
Daily Aspirin Use		Expected	98.6%
Statin Use		Above Expected	96.6%
Tobacco-Free		Expected	78.8%
Optimal Vascular Care		Expected	56.0%
BP Control		Expected	83.5%
Daily Aspirin Use		Expected	93.4%
Statin Use		Expected	95.6%
Tobacco-Free		Expected	74.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).