

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	86.0%
	Follow-up at Six Months	Expected	43.3%
	Response at Six Months	Expected	30.0%
	Remission at Six Months	Expected	13.3%
	Follow-up at 12 Months	Expected	53.3%
	Response at 12 Months	Expected	30.0%
	Remission at 12 Months	Expected	20.0%
Adolescent Mental Health and/or Depression Screening*		Average	91.8%
Colorectal Cancer Screening		Above Expected	81.1%
Optimal Asthma Control - Adults		Expected	55.0%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	63.4%
HbA1c Control		Expected	82.0%
BP Control		Expected	82.6%
Daily Aspirin Use		Expected	100.0%
Statin Use		Expected	93.8%
Tobacco-Free		Expected	92.6%
Optimal Vascular Care		Expected	63.9%
BP Control		Expected	83.3%
Daily Aspirin Use		Expected	88.9%
Statin Use		Expected	94.4%
Tobacco-Free		Expected	88.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).