

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	86.5%
	Follow-up at Six Months	Expected	50.0%
	Response at Six Months	Expected	13.6%
	Remission at Six Months	Expected	6.8%
	Follow-up at 12 Months	Expected	47.7%
	Response at 12 Months	Expected	11.4%
	Remission at 12 Months	Expected	4.6%
Adult Depression	PHQ-9 Utilization*	Above	90.4%
	Follow-up at Six Months	Above Expected	59.5%
	Response at Six Months	Expected	24.6%
	Remission at Six Months	Expected	10.9%
	Follow-up at 12 Months	Above Expected	58.5%
	Response at 12 Months	Expected	24.0%
	Remission at 12 Months	Expected	13.1%
Adolescent Mental Health and/or Depression Screening*		Below	88.3%
Colorectal Cancer Screening		Below Expected	69.7%
Optimal Asthma Control - Adults		Above Expected	65.4%
Optimal Asthma Control - Children		Above Expected	64.9%
Optimal Diabetes Care		Above Expected	50.5%
HbA1c Control		Expected	75.4%
BP Control		Expected	79.5%
Daily Aspirin Use		Expected	99.0%
Statin Use		Expected	87.3%
Tobacco-Free		Above Expected	91.3%
Optimal Vascular Care		Above Expected	63.0%
BP Control		Expected	84.4%
Daily Aspirin Use		Expected	86.1%
Statin Use		Expected	93.0%
Tobacco-Free		Above Expected	89.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).