

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	73.5%
	Follow-up at Six Months	Below Expected	27.8%
	Response at Six Months	Expected	15.3%
	Remission at Six Months	Expected	11.1%
	Follow-up at 12 Months	Below Expected	29.2%
	Response at 12 Months	Expected	15.3%
	Remission at 12 Months	Expected	8.3%
Adolescent Mental Health and/or Depression Screening*		Average	96.4%
Colorectal Cancer Screening		Expected	69.6%
Optimal Asthma Control - Adults		Below Expected	4.7%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	34.1%
HbA1c Control		Expected	68.8%
BP Control		Expected	78.9%
Daily Aspirin Use		Expected	98.2%
Statin Use		Below Expected	75.7%
Tobacco-Free		Expected	83.7%
Optimal Vascular Care		Expected	49.7%
BP Control		Expected	73.1%
Daily Aspirin Use		Expected	90.1%
Statin Use		Expected	89.4%
Tobacco-Free		Expected	80.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).