

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	81.9%
	Follow-up at Six Months	Below Expected	30.2%
	Response at Six Months	Below Expected	7.2%
	Remission at Six Months	Below Expected	3.6%
	Follow-up at 12 Months	Below Expected	32.4%
	Response at 12 Months	Expected	14.4%
	Remission at 12 Months	Expected	7.2%
Adolescent Mental Health and/or Depression Screening*		Average	93.3%
Colorectal Cancer Screening		Above Expected	74.6%
Optimal Asthma Control - Adults		Below Expected	18.8%
Optimal Asthma Control - Children		Below Expected	17.7%
Optimal Diabetes Care		Above Expected	53.2%
HbA1c Control		Expected	77.9%
BP Control		Above Expected	87.1%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	88.8%
Tobacco-Free		Expected	86.6%
Optimal Vascular Care		Expected	55.2%
BP Control		Above Expected	87.8%
Daily Aspirin Use		Below Expected	81.8%
Statin Use		Expected	89.6%
Tobacco-Free		Expected	81.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).