

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	93.2%
	Follow-up at Six Months	Expected	48.9%
	Response at Six Months	Expected	17.8%
	Remission at Six Months	Expected	10.5%
	Follow-up at 12 Months	Expected	54.3%
	Response at 12 Months	Expected	17.8%
	Remission at 12 Months	Expected	11.8%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Above Expected	79.5%
Optimal Asthma Control - Adults		Above Expected	80.1%
Optimal Asthma Control - Children		Expected	61.8%
Optimal Diabetes Care		Expected	52.5%
HbA1c Control		Expected	73.8%
BP Control		Above Expected	88.0%
Daily Aspirin Use		Expected	98.1%
Statin Use		Above Expected	93.0%
Tobacco-Free		Expected	87.6%
Optimal Vascular Care		Expected	62.9%
BP Control		Above Expected	87.4%
Daily Aspirin Use		Expected	84.6%
Statin Use		Expected	94.0%
Tobacco-Free		Expected	87.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).