

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	64.3%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	90.3%
Colorectal Cancer Screening		Below Expected	57.2%
Optimal Asthma Control - Adults		Below Expected	12.5%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	34.5%
HbA1c Control		Expected	64.5%
BP Control		Above Expected	90.9%
Daily Aspirin Use		Expected	98.5%
Statin Use		Below Expected	78.2%
Tobacco-Free		Below Expected	68.5%
Optimal Vascular Care		Expected	42.9%
BP Control		Expected	88.3%
Daily Aspirin Use		Expected	76.6%
Statin Use		Below Expected	77.9%
Tobacco-Free		Expected	70.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).