

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	39.5%
	Follow-up at Six Months	Below Expected	19.7%
	Response at Six Months	Below Expected	2.8%
	Remission at Six Months	Below Expected	0.6%
	Follow-up at 12 Months	Expected	39.9%
	Response at 12 Months	Below Expected	10.7%
	Remission at 12 Months	Expected	6.2%
Adolescent Mental Health and/or Depression Screening*		Below	50.9%
Colorectal Cancer Screening		Above Expected	73.3%
Optimal Asthma Control - Adults		Below Expected	17.3%
Optimal Asthma Control - Children		Below Expected	20.3%
Optimal Diabetes Care		Below Expected	36.4%
HbA1c Control		Expected	71.0%
BP Control		Expected	81.4%
Daily Aspirin Use		Below Expected	94.0%
Statin Use		Below Expected	80.1%
Tobacco-Free		Below Expected	81.5%
Optimal Vascular Care		Below Expected	37.8%
BP Control		Expected	82.6%
Daily Aspirin Use		Below Expected	66.9%
Statin Use		Below Expected	83.6%
Tobacco-Free		Below Expected	75.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).