

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	80.5%
	Follow-up at Six Months	Below Expected	27.0%
	Response at Six Months	Expected	9.5%
	Remission at Six Months	Expected	5.4%
	Follow-up at 12 Months	Expected	33.8%
	Response at 12 Months	Expected	16.2%
	Remission at 12 Months	Expected	8.1%
Adolescent Mental Health and/or Depression Screening*		Below	84.5%
Colorectal Cancer Screening		Below Expected	55.6%
Optimal Asthma Control - Adults		Above Expected	54.4%
Optimal Asthma Control - Children		Above Expected	59.1%
Optimal Diabetes Care		Below Expected	35.2%
HbA1c Control		Expected	66.4%
BP Control		Below Expected	71.1%
Daily Aspirin Use		Expected	98.3%
Statin Use		Above Expected	89.7%
Tobacco-Free		Expected	81.1%
Optimal Vascular Care		Below Expected	40.3%
BP Control		Below Expected	67.9%
Daily Aspirin Use		Expected	83.7%
Statin Use		Expected	90.1%
Tobacco-Free		Expected	70.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).