

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	67.4%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Expected	43.3%
Optimal Asthma Control - Adults		Expected	35.4%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	14.7%
HbA1c Control		Below Expected	48.6%
BP Control		Expected	74.3%
Daily Aspirin Use		Expected	98.2%
Statin Use		Expected	91.7%
Tobacco-Free		Below Expected	39.5%
Optimal Vascular Care		Expected	25.9%
BP Control		Expected	79.6%
Daily Aspirin Use		Expected	94.4%
Statin Use		Expected	94.4%
Tobacco-Free		Below Expected	37.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).