

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	81.3%
	Follow-up at Six Months	Expected	43.5%
	Response at Six Months	Expected	12.3%
	Remission at Six Months	Expected	7.8%
	Follow-up at 12 Months	Expected	40.9%
	Response at 12 Months	Expected	14.3%
	Remission at 12 Months	Expected	8.4%
Adolescent Mental Health and/or Depression Screening*		Average	96.1%
Colorectal Cancer Screening		Below Expected	63.0%
Optimal Asthma Control - Adults		Expected	50.0%
Optimal Asthma Control - Children		Expected	32.6%
Optimal Diabetes Care		Expected	40.4%
HbA1c Control		Expected	72.1%
BP Control		Expected	79.3%
Daily Aspirin Use		Expected	99.3%
Statin Use		Expected	86.5%
Tobacco-Free		Below Expected	76.9%
Optimal Vascular Care		Below Expected	43.7%
BP Control		Expected	74.2%
Daily Aspirin Use		Expected	90.6%
Statin Use		Expected	91.1%
Tobacco-Free		Below Expected	70.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).