

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	55.3%
	Follow-up at Six Months	Expected	40.9%
	Response at Six Months	Expected	15.5%
	Remission at Six Months	Expected	7.0%
	Follow-up at 12 Months	Expected	36.6%
	Response at 12 Months	Expected	8.5%
	Remission at 12 Months	Below Expected	1.4%
Adolescent Mental Health and/or Depression Screening*		Average	87.9%
Colorectal Cancer Screening		Below Expected	52.0%
Optimal Asthma Control - Adults		Expected	50.0%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	20.0%
HbA1c Control		Below Expected	54.0%
BP Control		Expected	77.3%
Daily Aspirin Use		Expected	100.0%
Statin Use		Below Expected	72.7%
Tobacco-Free		Below Expected	72.7%
Optimal Vascular Care		Expected	44.8%
BP Control		Expected	75.9%
Daily Aspirin Use		Above Expected	98.3%
Statin Use		Expected	89.7%
Tobacco-Free		Expected	70.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).