

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Below	45.5%
	Follow-up at Six Months	Above Expected	63.3%
	Response at Six Months	Expected	13.3%
	Remission at Six Months	Expected	4.4%
	Follow-up at 12 Months	Expected	53.3%
	Response at 12 Months	Expected	16.7%
	Remission at 12 Months	Expected	8.9%
Adult Depression	PHQ-9 Utilization*	Above	84.5%
	Follow-up at Six Months	Above Expected	60.1%
	Response at Six Months	Expected	16.0%
	Remission at Six Months	Expected	7.9%
	Follow-up at 12 Months	Expected	49.7%
	Response at 12 Months	Expected	14.6%
	Remission at 12 Months	Expected	7.3%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		-	-
HbA1c Control		-	-
BP Control		-	-
Daily Aspirin Use		-	-
Statin Use		-	-
Tobacco-Free		-	-
Optimal Vascular Care		-	-
BP Control		-	-
Daily Aspirin Use		-	-
Statin Use		-	-
Tobacco-Free		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).