

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	83.9%
	Follow-up at Six Months	Expected	52.0%
	Response at Six Months	Below Expected	10.9%
	Remission at Six Months	Below Expected	4.5%
	Follow-up at 12 Months	Expected	54.5%
	Response at 12 Months	Expected	14.4%
	Remission at 12 Months	Expected	6.4%
Adolescent Mental Health and/or Depression Screening*		Average	98.3%
Colorectal Cancer Screening		Above Expected	72.4%
Optimal Asthma Control - Adults		Above Expected	66.5%
Optimal Asthma Control - Children		Expected	64.8%
Optimal Diabetes Care		Expected	39.4%
HbA1c Control		Expected	68.9%
BP Control		Expected	75.4%
Daily Aspirin Use		Expected	98.7%
Statin Use		Expected	88.6%
Tobacco-Free		Expected	82.5%
Optimal Vascular Care		Expected	54.5%
BP Control		Expected	77.6%
Daily Aspirin Use		Expected	88.2%
Statin Use		Expected	91.5%
Tobacco-Free		Expected	80.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).