

**Clinical Quality Measures**

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	96.0%
Colorectal Cancer Screening		Below Expected	67.3%
Optimal Asthma Control - Adults		Below Expected	29.7%
Optimal Asthma Control - Children		Below Expected	14.0%
Optimal Diabetes Care		Below Expected	39.9%
HbA1c Control		Below Expected	68.6%
BP Control		Expected	82.0%
Daily Aspirin Use		Expected	99.0%
Statin Use		Expected	87.9%
Tobacco-Free		Below Expected	80.4%
Optimal Vascular Care		Expected	62.2%
BP Control		Expected	83.9%
Daily Aspirin Use		Expected	90.6%
Statin Use		Expected	93.9%
Tobacco-Free		Expected	77.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

**Ratings**

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).