

Clinical Quality Measures

2023 measurement year

| Measure | | Rating | Rate |
|---|-------------------------|----------------|-------|
| Adolescent Depression | PHQ-9 Utilization* | - | - |
| | Follow-up at Six Months | - | - |
| | Response at Six Months | - | - |
| | Remission at Six Months | - | - |
| | Follow-up at 12 Months | - | - |
| | Response at 12 Months | - | - |
| | Remission at 12 Months | - | - |
| Adult Depression | PHQ-9 Utilization* | - | - |
| | Follow-up at Six Months | - | - |
| | Response at Six Months | - | - |
| | Remission at Six Months | - | - |
| | Follow-up at 12 Months | - | - |
| | Response at 12 Months | - | - |
| | Remission at 12 Months | - | - |
| Adolescent Mental Health and/or Depression Screening* | | Above | 99.3% |
| Colorectal Cancer Screening | | Expected | 70.0% |
| Optimal Asthma Control - Adults | | Expected | 54.6% |
| Optimal Asthma Control - Children | | Above Expected | 66.7% |
| Optimal Diabetes Care | | Expected | 54.5% |
| HbA1c Control | | Above Expected | 81.8% |
| BP Control | | Above Expected | 86.5% |
| Daily Aspirin Use | | Expected | 99.4% |
| Statin Use | | Expected | 88.9% |
| Tobacco-Free | | Expected | 83.8% |
| Optimal Vascular Care | | Expected | 57.3% |
| BP Control | | Expected | 81.9% |
| Daily Aspirin Use | | Above Expected | 91.3% |
| Statin Use | | Expected | 94.9% |
| Tobacco-Free | | Expected | 77.5% |

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).