

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	98.1%
Colorectal Cancer Screening		Below Expected	67.8%
Optimal Asthma Control - Adults		Below Expected	30.5%
Optimal Asthma Control - Children		Expected	53.2%
Optimal Diabetes Care		Below Expected	39.6%
HbA1c Control		Below Expected	69.5%
BP Control		Expected	81.2%
Daily Aspirin Use		Above Expected	99.2%
Statin Use		Below Expected	83.9%
Tobacco-Free		Below Expected	81.1%
Optimal Vascular Care		Expected	52.6%
BP Control		Expected	79.4%
Daily Aspirin Use		Above Expected	91.6%
Statin Use		Expected	91.8%
Tobacco-Free		Below Expected	77.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).