

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	81.2%
	Follow-up at Six Months	Expected	43.8%
	Response at Six Months	Expected	20.6%
	Remission at Six Months	Expected	11.0%
	Follow-up at 12 Months	Expected	56.2%
	Response at 12 Months	Expected	27.4%
	Remission at 12 Months	Expected	20.6%
Adolescent Mental Health and/or Depression Screening*		Average	96.2%
Colorectal Cancer Screening		Above Expected	76.3%
Optimal Asthma Control - Adults		Expected	50.0%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	55.3%
HbA1c Control		Expected	77.4%
BP Control		Expected	85.7%
Daily Aspirin Use		Expected	100.0%
Statin Use		Above Expected	94.0%
Tobacco-Free		Expected	83.0%
Optimal Vascular Care		Expected	57.3%
BP Control		Expected	77.4%
Daily Aspirin Use		Above Expected	96.8%
Statin Use		Expected	95.2%
Tobacco-Free		Expected	79.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).